

July, 2002
hep-ph/0207226

The contribution of off-shell gluons to the longitudinal structure function F_L

A.V. Kotikov

*Bogoliubov Laboratory of Theoretical Physics
Joint Institute for Nuclear Research
141980 Dubna, Russia*

A.V. Lipatov

*Department of Physics
Lomonosov Moscow State University
119899 Moscow, Russia*

N.P. Zotov

*Skobeltsyn Institute of Nuclear Physics
Lomonosov Moscow State University
119992 Moscow, Russia*

Abstract

We present results for the structure function F_L for a gluon target having nonzero transverse momentum square at order α_s . The results of double convolution (with respect to Bjorken variable x and the transverse momentum) of the perturbative part and the unintegrated gluon densities are compared with recent experimental data for F_L at low x values and with predictions of other approaches.

PACS number(s): 13.60.Hb, 12.38.Bx, 13.15.Dk

1 Introduction

The basic information on the internal structure of nucleons is extracted from the process of deep inelastic (lepton-hadron) scattering (DIS). Its differential cross-section has the form:

$$\frac{d^2\sigma}{dx dy} = \frac{2\pi\alpha_{em}^2}{xQ^4} [(1 - y + y^2/2) F_2(x, Q^2) - (y^2/2) F_L(x, Q^2)],$$

where $F_2(x, Q^2)$ and $F_L(x, Q^2)$ are the transverse and longitudinal structure functions (SF), respectively, q^μ and p^μ are the photon and the hadron 4-momentums and $x = Q^2/(2pq)$ with $Q^2 = -q^2 > 0$.

The longitudinal SF $F_L(x, Q^2)$ is a very sensitive QCD characteristic because it is equal to zero in the parton model with spin-1/2 partons. Unfortunately, essentially at small values of x , the experimental extraction of F_L data is required a rather cumbersome procedure (see [1, 2], for example). Moreover, the perturbative QCD leads to some controversial results in the case of SF F_L . The next-to-leading order (NLO) corrections to the longitudinal coefficient function, which are large and negative at small x [3, 4], need a resummation procedure ¹ that leads to coupling constant scale higher essentially then Q^2 (see [6, 4, 7]) ².

Recently there have been important new data [13]-[18] of the longitudinal SF F_L , which have probed the small- x region down to $x \sim 10^{-2}$. Moreover, the SF F_L can be related at small x with SF F_2 and the derivation $dF_2/d\ln(Q^2)$ (see [19]-[21]). In this way most precise predictions based on data of F_2 and $dF_2/d\ln(Q^2)$ (see [15] and references therein) can be obtained for F_L . These predictions can be considered as indirect 'experimental data' for F_L .

In this paper for analysis of the above data we will use so called k_T -factorization approach [22, 23, 24] based on BFKL dynamics [12] (see also recent review [11] and references therein). In the framework of the k_T -factorization approach, a study of the longitudinal SF F_L has been done firstly in Ref. [25], where small x asymptotics of F_L has been evaluated using the BFKL results for the Mellin transform of the unintegrated gluon distribution and the longitudinal Wilson coefficient functions has been calculated analytically for the full perturbative series at asymptotically small x values. Since we want to analyze F_L data in a broader range at small x , we will use parameterizations of the unintegrated gluon distribution function $\Phi_g(x, k_\perp^2)$ (see Section 3).

The unintegrated gluon distribution $\Phi_g(x, k_\perp^2)$ (f_g is the (integrated) gluon distribution in proton multiplied by x and $k_\perp$ is the transverse part of gluon 4-momentum k^μ)

$$f_g(x, Q^2) = \int^{Q^2} dk_\perp^2 \Phi_g(x, k_\perp^2) \quad (\text{hereafter } k^2 = -k_\perp^2) \quad (1)$$

is the basic dynamical quantity in the k_T -factorization approach ³. It satisfies the BFKL equation [12].

¹ Without a resummation the NLO approximation of SF F_L can be negative at low x and quite low Q^2 values (see [4, 5]).

² Note that at low x a similar property has been observed also in the approaches [8, 9, 10] (see recent review [11] and discussions therein), which based on Balitsky-Fadin-Kuraev-Lipatov (BFKL) dynamics [12], where the leading $\ln(1/x)$ contributions are summed.

³ In our previous analysis [26] we shown that the property $k^2 = -k_\perp^2$ leads to the equality of the Bjorken x value in the standard renormalization-group approach and in the Sudakov one.

Notice that the integral is divergent at lower limit (at least, for some parameterizations of $\Phi_g(x, k_\perp^2)$) and so it leads to the necessity to consider the difference $f_g(x, Q^2) - f_g(x, Q_0^2)$ with some nonzero Q_0^2 (see discussions in [26]), i.e.

$$f_g(x, Q^2) = f_g(x, Q_0^2) + \int_{Q_0^2}^{Q^2} dk_\perp^2 \Phi_g(x, k_\perp^2) \quad (2)$$

Then, in the k_T -factorization the SF $F_{2,L}(x, Q^2)$ are driven at small x primarily by gluons and are related in the following way to the unintegrated distribution $\Phi_g(x, k_\perp^2)$:

$$F_{2,L}(x, Q^2) = \int_x^1 \frac{dz}{z} \int dk_\perp^2 \sum_{i=u,d,s,c} e_i^2 \cdot \hat{C}_{2,L}^g(x/z, Q^2, m_i^2, k_\perp^2) \Phi_g(z, k_\perp^2), \quad (3)$$

where e_i^2 are charge squares of active quarks.

The functions $\hat{C}_{2,L}^g(x, Q^2, m_i^2, k_\perp^2)$ can be regarded as SF of the off-shell gluons with virtuality $k_\perp^2$ (hereafter we call them as *hard structure functions*⁴). They are described by the sum of the quark boxes (and crossed boxes) diagram contribution to the photon-gluon interaction (see Fig. 1).

The purpose of the paper is to give predictions for the longitudinal SF $F_L(x, Q^2)$ based on the calculations of the hard SF $\hat{C}_{2,L}^g(x, Q^2, m^2, k_\perp^2)$, given in our previous study [26], and several parameterizations of unintegrated gluon distributions (see [11] and references therein).

It is instructive to note that the diagrams shown in Fig. 1. are similar to those of the photon-photon scattering process. The corresponding QED contributions have been calculated many years ago in Ref. [27] (see also the beautiful review in Ref. [28]). Our results have been calculated independently and they are in full agreement with Ref. [27]. Moreover, our results are in agreement with the corresponding integral representations for $\hat{C}_{2,L}^g$, given in [22, 25] and numerically with results of [29]. However, we hope that our formulas which are given in a simpler form could be useful for others. This simpler form for the hard SF $\hat{C}_{2,L}^g$ comes from using the relation between results based on non-sense, transverse and longitudinal gluon polarizations (see Eq. (13) below) observed in [26] for gauge-invariant sets of diagrams.

The structure of this paper is as follows: in Section 2 we present the basic formulae of our approach. Section 3 contains the relations between SF F_L and F_2 and the derivative $dF_2/d\ln Q^2$, obtained in [19, 20, 21] in the framework of Dokshitzer-Gribov-Lipatov-Altarelli-Parisi (DGLAP) approach [30] (i.e. in the collinear approximation: $k_\perp^2 = 0$). In Section 4 we give the predictions for the structure function F_L for three cases of unintegrated gluon distributions.

2 Basic formulae

To begin with, we review shortly results of Ref. [26] needed below in our investigations.

The hadron part of the DIS spin-average lepton-hadron cross section can be represented in the form⁵

$$F_{\mu\nu} = e_{\mu\nu}(q) F_L(x, Q^2) + d_{\mu\nu}(q, p) F_2(x, Q^2), \quad (4)$$

⁴by analogy with similar relations between cross-sections and hard cross-sections.

⁵Hereafter we consider only one-photon exchange approximation.

where

$$e_{\mu\nu}(q) = g_{\mu\nu} - \frac{q_\mu q_\nu}{q^2} \quad \text{and} \quad d_{\mu\nu}(q, p) = - \left[g_{\mu\nu} + 2x_B \frac{(p_\mu q_\nu + p_\nu q_\mu)}{q^2} + p_\mu p_\nu \frac{4x_B^2}{q^2} \right]$$

2.1 Feynman-gauge gluon polarization

As it has been shown in Ref. [26], it is very convenient to consider, as the first approximation, gluons having polarization tensor (hereafter the indices α and β are connected with gluons and μ and ν are connected with photons)⁶:

$$\hat{P}^{\alpha\beta} = -g^{\alpha\beta} \quad (5)$$

The tensor corresponds is equal to the standard choice of polarization matrix in the framework of collinear approximation. In a sense the case of polarization is equal to the standard DIS suggestions about parton properties, excepting their off-shell property. The polarization (5) gives the main contribution to the polarization tensor we are interested in (see below)

$$\hat{P}_{BFKL}^{\alpha\beta} = \frac{k_\perp^\alpha k_\perp^\beta}{k_\perp^2}, \quad (6)$$

which comes from the high energy (or k_T) factorization prescription [22, 23, 24]⁷.

Contracting the photon projectors (connected with photon indices of diagrams on Fig.1.)

$$\hat{P}_{\mu\nu}^{(1)} = -\frac{1}{2}g_{\mu\nu} \quad \text{and} \quad \hat{P}_{\mu\nu}^{(2)} = 4z^2 \frac{k_\mu k_\nu}{Q^2},$$

(here $z = Q^2/(2pq)$ is the corresponding Bjorken variable on parton level) with the parton tensor $F_{\mu\nu}^p$:

$$F_{\mu\nu}^p = e_{\mu\nu}(q) F_L^p(z, Q^2) + d_{\mu\nu}(q, k) F_2^p(z, Q^2), \quad (7)$$

we obtain at the parton level (i.e. for off-shell gluons having momentum k_μ), when $\hat{C}_{2,L}^g(z) \sim F_{2,L}^p(z, Q^2)$ ⁸:

$$\tilde{\beta}^2 \cdot \hat{C}_2^g(x) = \mathcal{K} \cdot \left[f^{(1)} + \frac{3}{2\tilde{\beta}^2} \cdot f^{(2)} \right] \quad (8)$$

$$\tilde{\beta}^2 \cdot \hat{C}_L^g(x) = \mathcal{K} \cdot \left[4bx^2 f^{(1)} + \frac{(1 + 2bx^2)}{\tilde{\beta}^2} \cdot f^{(2)} \right] = \mathcal{K} \cdot f^{(2)} + 4bx^2 \tilde{\beta}^2 \cdot C_2^g, \quad (9)$$

⁶In principle, we can use here more general cases of polarization tensor (for example, one is based on Landau or unitary gauge). The difference between them and (5) is $\sim k^\alpha$ and/or $\sim k^\beta$ and, hence, it leads to zero contributions because Feynman diagrams on Fig.1 are gauge invariant.

⁷We would like to note that the BFKL-like polarization tensor (6) is a particular case of so-called nonsense polarization of the particles in t -channel. The nonsense polarization makes the main contributions for cross sections in s -channel at $s \rightarrow \infty$ (see, for example, Ref. [31] and references therein). The limit $s \rightarrow \infty$ corresponds to the small values of Bjorken variable x , that is just the range of our study.

⁸ The hard SF $\hat{C}_{2,L}^g$ do not depend on the type of target, so we can replace $z \rightarrow x$ below.

where the normalization factor $\mathcal{K} = a_s(Q^2) \cdot x$,

$$\hat{P}_{\mu\nu}^{(i)} F_{\mu\nu} = \mathcal{K} \cdot f^{(i)}, \quad i = 1, 2$$

and $a_s(Q^2) = \alpha_s(Q^2)/(4\pi)$, $\tilde{\beta}^2 = 1 - 4bx^2$, $b = -k^2/Q^2 \equiv k_\perp^2/Q^2 > 0$, $a = m^2/Q^2$.

Applying the projectors $\hat{P}_{\mu\nu}^{(i)}$ to Feynman diagrams displayed in Fig.1, we obtain the following results

$$\begin{aligned} f^{(1)} &= -2\beta \left[1 - \left(1 - 2x(1+b-2a) \cdot [1 - x(1+b+2a)] \right) \cdot f_1 \right. \\ &\quad \left. + (2a-b)(1-2a)x^2 \cdot f_2 \right], \end{aligned} \quad (10)$$

$$\begin{aligned} f^{(2)} &= 8x \cdot \beta \left[(1 - (1+b)x) - 2x \left(bx(1 - (1+b)x)(1+b-2a) + a\tilde{\beta}^2 \right) \cdot f_1 \right. \\ &\quad \left. + bx^2(1 - (1+b)x)(2a-b) \cdot f_2 \right], \end{aligned} \quad (11)$$

where

$$\beta^2 = 1 - \frac{4ax}{(1 - (1+b)x)} \quad (12)$$

and

$$f_1 = \frac{1}{\tilde{\beta}\beta} \cdot \ln \frac{1 + \beta\tilde{\beta}}{1 - \beta\tilde{\beta}}, \quad f_2 = -\frac{4}{1 - \beta^2\tilde{\beta}^2}$$

2.2 BFKL-like gluon polarization

Now we take into account the BFKL-like gluon polarization (6). As we shown in [26], the projector $\hat{P}_{BFKL}^{\alpha\beta}$ can be represented as

$$\hat{P}_{BFKL}^{\alpha\beta} = -\frac{1}{2} \frac{1}{\tilde{\beta}^4} \left[\tilde{\beta}^2 g^{\alpha\beta} - 12bx^2 \frac{q^\alpha q^\beta}{Q^2} \right] \quad (13)$$

In the previous subsection we have already presented the contributions to hard SF using the first term in the brackets of the r.h.s. of (13). Repeating the above calculations with the projector $\sim q^\alpha q^\beta$, we obtain the total contributions to hard SF which can be represented as the following shift of the results in Eqs.(8) - (11):

$$\begin{aligned} \hat{C}_2^g(x) &\rightarrow \hat{C}_{2,BFKL}^g(x), \quad \hat{C}_L^g(x) \rightarrow \hat{C}_{L,BFKL}^g(x); \\ f^{(1)} &\rightarrow f_{BFKL}^{(1)} = \frac{1}{\tilde{\beta}^4} \left[\tilde{\beta}^2 f^{(1)} - 3bx^2 \tilde{f}^{(1)} \right] \\ f^{(2)} &\rightarrow f_{BFKL}^{(2)} = \frac{1}{\tilde{\beta}^4} \left[\tilde{\beta}^2 f^{(2)} - 3bx^2 \tilde{f}^{(2)} \right], \end{aligned} \quad (14)$$

where

$$\begin{aligned}\tilde{f}^{(1)} &= -\beta \left[\frac{1-x(1+b)}{x} - 2 \left(x(1-x(1+b))(1+b-2a) + a\tilde{\beta}^2 \right) \cdot f_1 \right. \\ &\quad \left. - x(1-x(1+b))(1-2a) \cdot f_2 \right],\end{aligned}\tag{15}$$

$$\tilde{f}^{(2)} = 4 \cdot \beta (1 - (1+b)x)^2 \left[2 - (1+2bx^2) \cdot f_1 - bx^2 \cdot f_2 \right],\tag{16}$$

Notice that the general formulae are needed only to evaluate the charm contribution to structure functions F_2 and F_L , i.e. F_2^c and F_L^c . To evaluate the corresponding light-quark contributions, i.e. F_L^l , we can use $m^2 = 0$ limit of above formulae.

2.3 The case $m^2 = 0$

When $m^2 = 0$ the hard SF $\hat{C}_k^g(x)$ are defined by $f^{(1)}$, $f^{(2)}$, $\tilde{f}^{(1)}$ and $\tilde{f}^{(2)}$ (as in Eqs.(8), (9) and (14)) which can be represented as

$$f^{(1)} = -2 \left[2 - \left(1 - 2x(1+b) + 2x^2(1+b)^2 \right) \cdot L(\tilde{\beta}) \right],\tag{17}$$

$$f^{(2)} = 8x(1+b)(1-(1+b)x) \left[1 - 2bx^2 \cdot L(\tilde{\beta}) \right],\tag{18}$$

$$\tilde{f}^{(1)} = -\frac{(1+b)(1-x(1+b))}{bx} \left[1 - 2bx^2 \cdot L(\tilde{\beta}) \right] = -\frac{1}{8bx^2} f^{(2)},\tag{19}$$

$$\tilde{f}^{(2)} = 4(1-x(1+b))^2 \left[3 - (1+2bx^2) \cdot L(\tilde{\beta}) \right]\tag{20}$$

and, thus, (see Eq. (14))

$$\begin{aligned}\tilde{\beta}^4 \cdot f_{BFKL}^{(1)} &= (-2)(1-x(1+b)) \left[2 \left(1 - 2x(1+b) + \frac{x^2(1-b)^2}{1-x(1+b)} \right) \right. \\ &\quad \left. - \left(1 - x(1+b) - 4x^3b(1+b) + \frac{x^2(1-b)^2}{1-x(1+b)} \right) \cdot L(\tilde{\beta}) \right],\end{aligned}\tag{21}$$

$$\begin{aligned}\tilde{\beta}^4 \cdot f_{BFKL}^{(2)} &= 8x(1-x(1+b)) \left[1 + b - 18bx(1-x(1+b)) \right. \\ &\quad \left. + 2bx \left(3 - 4x(1+b) + 6bx^2(1-x(1+b)) \right) \cdot L(\tilde{\beta}) \right],\end{aligned}\tag{22}$$

where

$$L(\tilde{\beta}) = \frac{1}{\tilde{\beta}} \cdot \ln \frac{1+\tilde{\beta}}{1-\tilde{\beta}}$$

3 Relations between F_L , F_2 and derivation of F_2 in the case of collinear approximation

Another information about the SF F_L can be obtained in the collinear approximation (i.e. when $k_\perp^2 = 0$) in the following way.

In the framework of perturbative QCD, there is the possibility to connecting F_L to F_2 and its derivation $dF_2/d\ln Q^2$ due the fact that at small x the DIS structure functions depend only on two *independent* functions: the gluon distribution and singlet quark one (the nonsinglet quark density is negligible at small x), which in turn can be expressed in terms of measurable SF F_2 and its derivation $dF_2/d\ln Q^2$.

In this way, by analogy with the case of the gluon distribution function (see [32, 33] and references therein), the behavior of $F_L(x, Q^2)$ has been studied in [19]-[21], using the HERA data [34, 35] and the method [36]⁹ of replacement of the Mellin convolution by ordinary products. Thus, the small x behavior of the SF $F_L(x, Q^2)$ can be extracted directly from the measured values of $F_2(x, Q^2)$ and its derivative without a cumbersome procedure (see [1, 2]). These extracted values of F_L may be well considered as *new small x 'experimental data' of F_L* . The relations can be violated by nonperturbative corrections like higher twist ones (see [38, 39]), which can be large exactly in the case of SF F_L [25, 40].

Because k_T -factorization approach is one of popular nonperturbative approaches used at small x , it is very useful to compare its predictions with the results of [19]-[21] based on the relations between SF $F_L(x, Q^2)$, $F_2(x, Q^2)$ and $dF_2(x, Q^2)/d\ln Q^2$. It is the main purpose of the study.

The k_T -factorization approach relates strongly to Regge-like behavior of parton distributions. So, we restrict our investigations to SF and parton distributions at the following form (hereafter $a = q, g$):

$$f_a(x, Q^2) \sim F_2(x, Q^2) \sim x^{-\delta(Q^2)} \quad (23)$$

Note that really the slopes of the sea quark and gluon distributions: δ_q and δ_g , respectively, and the slope δ_{F_2} of SF F_2 are little different. The slopes have a familiar property $\delta_q < \delta_{F_2} < \delta_g$ (see Refs. [41]-[46] and references therein). We will neglect, however, this difference and use in our investigations the experimental values of $\delta(Q^2) \equiv \delta_{F_2}(Q^2)$ extracted by H1 Collaboration¹⁰ (see [41] and references therein). We note that the Q^2 -dependence is in very good agreement with perturbative QCD at $Q^2 \geq 2 \text{ GeV}^2$ (see [48]). Moreover, the values of the slope $\delta(Q^2)$ are in agreement with recent phenomenological studies (see, for example, [8]) incorporating the next-to-leading corrections [49] (see also [50]) in the framework of BFKL approach.

Thus, assuming the *Regge-like behavior* (23) for the gluon distribution and $F_2(x, Q^2)$ at $x^{-\delta} \gg 1$ and using the NLO approximation for collinear coefficient functions and anomalous

⁹The method is based on previous investigations [1, 37].

¹⁰ Now the preliminary ZEUS data for the slope $d\ln F_2/d\ln(1/x)$ are available as some points on Figs. 8 and 9 in Ref. [47]. Moreover, the new preliminary H1 points have been presented on the Workshop DIS2002 (see [17]). Both the new points are shown quite similar properties to compare with H1 data [41]. Unfortunately, tables of the ZEUS data and the new H1 data are unavailable yet and, so, the points cannot be used here.

dimensions of Wilson operators, the following results for $F_L(x, Q^2)$ has been obtained in [20]:

$$F_L(x, Q^2) = -2 \frac{B_L^{g,1+\delta} (1 + a_s(Q^2) \bar{R}_L^{g,1+\delta})}{\gamma_{qg}^{(0),1+\delta} + \bar{\gamma}_{qg}^{(1),1+\delta} a_s(Q^2)} \xi^\delta \left[\frac{dF_2(x\xi, Q^2)}{d \ln Q^2} \right. \\ \left. + \frac{a_s(Q^2)}{2} \left(\frac{B_L^{g,1+\delta}}{B_L^{g,1+\delta}} \gamma_{qg}^{(0),1+\delta} - \gamma_{qq}^{(0),1+\delta} \right) F_2(x\xi, Q^2) \right] + O(a_s^2, x^{2-\delta}, \alpha x^{1-\delta}), \quad (24)$$

where

$$\bar{\gamma}_{qg}^{(1),\eta} = \gamma_{qg}^{(1),\eta} + B_2^{q,\eta} \gamma_{qg}^{(0),\eta} + B_2^{g,\eta} (2\beta_0 + \gamma_{gg}^{(0),\eta} - \gamma_{qq}^{(0),\eta}), \quad \bar{R}_L^{g,\eta} = R_L^{g,\eta} - B_2^{g,\eta} \frac{B_L^{q,\eta}}{B_L^{g,\eta}}$$

Here $\bar{\gamma}_{qa}^{(1),\eta}$ and $\bar{R}_L^{a,\eta}$ ($a = q, g$) are the combinations¹¹ of the 'anomalous dimensions' of Wilson operators $\gamma_{qa}^\eta = a_s \gamma_{qa}^{(0),\eta} + a_s^2 \gamma_{qa}^{(1),\eta} + O(a_s^3)$ and 'Wilson coefficients' $a_s B_L^{a,\eta} (1 + a_s R_L^{a,\eta}) + O(a_s^3)$ and $a_s B_2^{a,\eta} + O(a_s^2)$ with the 'moment' argument η (i.e., the combinations of the functions which can be obtained by analytical continuation of the corresponding anomalous dimensions and coefficient functions from integer values n of their argument to non-integer ones η).

Note that, in principle, any term like $\sim 1/(n+m)$ ($m = 0, 1, 2, \dots$) which comes to the corresponding combinations of the anomalous dimensions and coefficient functions: $\bar{\gamma}_{qa}^{(1),n}$ and $\bar{R}_L^{a,n}$, should contribute to Eq.(24) in the following form (after replacement of Mellin convolutions by usual products in the DGLAP equations (see [36])):

$$\frac{1}{1+\delta+m} \left(1 + \frac{\Gamma(2+\delta+m)\Gamma(1+\nu)}{\Gamma(2+\delta+m+1+\nu)} \cdot x^{1+\delta+m} \right), \quad (25)$$

where the value of ν comes (see [51, 52, 53]) from asymptotics of parton distributions $f_a(x)$ at $x \rightarrow 1$: $f_a \sim (1-x)^{\nu_a}$, and¹² $\nu \approx 4$ from quark account rules [54]. The additional term $\sim x^{1+\delta+m}$ in Eq. (25) is important only for $m = -1$ case (i.e. for the singular parts $\sim 1/(n-1)$ of the corresponding anomalous dimensions and coefficient functions) and quite small values of δ (i.e. for $x^\delta \sim Const$).

Thus, excepting the case when $m = -1$ and $x^\delta \sim Const$, we can replace (25) by its first term $1/(1+\delta+m)$, i.e. our variables $\bar{\gamma}_{qg}^{(1),\eta}$ and $\bar{R}_L^{g,\eta}$ are just the combinations of corresponding anomalous dimensions and coefficient functions at $n = 1 + \delta$. When $m = -1$ and $x^\delta \sim Const$ at the considered small x range, we should replace the term $1/(n-1)$, if it was exist in the variables $\bar{\gamma}_{qa}^{(1),n}$ and $\bar{R}_L^{a,n}$ ($a = q, g$), by the following term:

$$\frac{1}{\tilde{\delta}} = \frac{1}{\delta} \left[1 - \frac{\Gamma(1-\delta)\Gamma(1+\nu)}{\Gamma(1-\delta+\nu)} x^\delta \right] \quad (26)$$

Note also that the $1/\tilde{\delta}$ coincides approximately with $1/\delta$ when $\delta \neq 0$ and $x \rightarrow 0$. However, at $\delta \rightarrow 0$, the value of $1/\tilde{\delta}$ is not singular:

$$\frac{1}{\tilde{\delta}} \rightarrow \ln \left(\frac{1}{x} \right) - \left[\Psi(1+\nu) - \Psi(1) \right] \quad (27)$$

¹¹Because we consider here $F_2(x, Q^2)$ but not the singlet quark distribution in the corresponding DGLAP equations [30].

¹²In our formula (24) we have our interest mostly to gluons, so we can apply $\nu = \nu_g \approx 4$ below.

So, the Eq. (24) together with well-known expressions of anomalous dimensions $\gamma_{ab}^{(0),n}$ and $\gamma_{ab}^{(1),n}$ ($a, b = q, g$) and coefficient functions $B_2^{a,n}$ and $R_L^{a,n}$ ($a = q, g$) (see [55] and [56, 57], respectively, and references therein) gives a possibility to extract SF F_L at small x values. The calculations are based on precise experimental data of SF F_2 and its derivatives $dF_2/d\ln(Q^2)$ and $\delta \equiv d\ln F_2/d\ln(1/x)$.

For concrete δ -values, the Eq. (24) simplifies essentially (see [20]). For example, for $\delta = 0.3$ we obtain (for the number of active quarks $f = 4$ and $\overline{MS}$ scheme):

$$F_L(x, Q^2) = \frac{0.84}{1 + 59.3a_s(Q^2)} \left[\frac{dF_2(0.48x, Q^2)}{d\ln Q^2} + 3.59a_s(Q^2)F_2(0.48x, Q^2) \right] + O(a_s^2 x^{2-\delta}, a_s x^{1-\delta}) \quad (28)$$

At arbitrary δ values, in real applications it is very useful to simplify Eq. (24) as follows. We keep the exact δ -dependence only for the leading order terms, which are very simple. In the NLO corrections we extract the terms $\sim 1/\tilde{\delta}$, which are changed strongly when $0 \leq \delta \leq 1$, and parameterize the rest terms in the form: $a_i + b_i\delta + c_i\delta^2$. The coefficients a_i , b_i , c_i are fixed from the agreement these parameterizations with the exact values of $\overline{\gamma}_{qa}^{(1),\eta}$ and $\overline{R}_L^{a,\eta}$ at $\delta = 0$, 0.3 and 0.5. These exact values can be found in Refs. [33, 20, 21].

Then, the approximate representation of Eq. (24) of for arbitrary δ value has the form:

$$F_L(x, Q^2) = \frac{r(1+\delta)(\xi(\delta))^\delta}{(1 + 30a_s(Q^2)[1/\tilde{\delta} - \frac{116}{45}\rho_1(\delta)])} \left[\frac{dF_2(x\xi(\delta), Q^2)}{d\ln Q^2} + \frac{8}{3}\rho_2(\delta)a_s(Q^2)F_2(x\xi(\delta), Q^2) \right] + O(a_s^2, a_s x, x^2), \quad (29)$$

where

$$\begin{aligned} r(\delta) &= \frac{4\delta}{2 + \delta + \delta^2}, & \xi(\delta) &= \frac{r(\delta)}{r(1+\delta)}, \\ \rho_1(\delta) &= 1 + \delta + \delta^2/4, & \rho_2(\delta) &= 1 - 2.39\delta + 2.69\delta^2, \end{aligned} \quad (30)$$

4 Comparison with F_L experimental data

With the help of the results obtained in the previous Section we have analyzed experimental data for SF F_L ¹³ from H1 [13] and [18], NMC [58], CCFR [59] and [60], BCDMS [61] collaborations ¹⁴. Note, that we do not correct CCFR data [59] and [60] which has been obtained in νN processes because the terms $\sim x \cdot m_c^2/Q^2$, which are different in μN and νN processes, are not so strong at low x .

We calculate the SF F_L as the sum of two types of contributions: charm quark one F_L^c and light quark one F_L^l :

$$F_L = F_L^l + F_L^c \quad (31)$$

¹³Sometimes there are experimental data for the ration $R = \sigma_L/\sigma_T$, which can be recalculated for the SF F_L because $F_L = F_2 R/(1 + R)$.

¹⁴ We do not use experimental data of R from SLAC [62, 63], EM [64] and CDHSW [65] Collaborations because they are obtained at quite large x values.

We use the expression (3) for the calculation of the both SF F_L^l and F_L^c in the following form (here $a_c = m_c^2/Q^2$):

$$F_L^l(x, Q^2) = \sum_{f=uds} e_f^2 \left[\int_x^1 \frac{dz}{z} \hat{C}_{L,BFKL}^g\left(\frac{x}{z}, Q^2, 0\right) z f_g(z, Q_0^2) + \sum_{i=1}^2 \int_{z_{min}^{(i)}}^{z_{max}^{(i)}} \frac{dz}{z} \int_{k_{\perp min}^{2(i)}}^{k_{\perp max}^{2(i)}} dk_{\perp}^2 \hat{C}_{L,BFKL}^g\left(\frac{x}{z}, Q^2, k_{\perp}^2\right) \Phi(z, k_{\perp}^2, Q_0^2) \right], \quad (32)$$

$$F_L^c(x, Q^2) = e_c^2 \left[\int_{x(1+4a_c)}^1 \frac{dz}{z} \hat{C}_{L,BFKL}^g\left(\frac{x}{z}, m_c^2, Q^2, 0\right) z f_g(z, Q_0^2) + \sum_{i=1}^2 \int_{z_{min}^{(i)}}^{z_{max}^{(i)}} \frac{dz}{z} \int_{k_{\perp min}^{2(i)}}^{k_{\perp max}^{2(i)}} dk_{\perp}^2 \hat{C}_{L,BFKL}^g\left(\frac{x}{z}, m_c^2, Q^2, k_{\perp}^2\right) \Phi(z, k_{\perp}^2, Q_0^2) \right], \quad (33)$$

where $\hat{C}_{L,BFKL}^g(x_B, Q^2, m_c^2, k_{\perp}^2)$ are given by Eqs. (9) and (14).

The integration limits in the expression (33) have the following values:

$$\begin{aligned} z_{min}^{(1)} &= x(1 + 4a_c + \frac{Q_0^2}{Q^2}), \quad z_{max}^{(1)} = 2x(1 + 2a_c); \\ k_{\perp min}^{2(1)} &= Q_0^2, \quad k_{\perp max}^{2(1)} = (\frac{z}{x} - (1 + 4a_c))Q^2; \\ z_{min}^{(2)} &= 2x(1 + 2a_c), \quad z_{max}^{(2)} = 1; \\ k_{\perp min}^{2(2)} &= Q_0^2, \quad k_{\perp max}^{2(2)} = Q^2; \end{aligned} \quad (34)$$

The ranges of integration correspond to positive values of square roots in expressions (10), (11), (15) and (16) and also should obey to kinematical restriction ($z \leq (1 + 4a_c + b)^{-1}$) following from condition $\beta^2 \geq 0$ (see Eqs. (10)-(12)). In Eq.(32) the ranges (34) are used at $a_c = 0$.

In Fig. 2 we show the SF F_L as a function x for different values of Q^2 in comparison with H1 experimental data sets: old one of [13] (black triangles), last year one of [15] (black squares) and new preliminary one of [18] (black circles) and also with NMC [58] (white triangles), CCFR [59] (white circles) and BCDMS [61] data (white squares). For comparison with these data we present the results of the calculation with three different parameterizations for the unintegrated gluon distribution $\Phi(x, k_{\perp}^2, Q_0^2)$ at $Q_0^2 = 4 \text{ GeV}^2$. All of them: Kwiecinski-Martin-Stasto (KMS) one [72] and Blumlein (JB) one [68] and Golec-Biernat and Wusthoff (GBW), have been used already in our previous work [26] and reviewed there.

There are several other popular parameterizations (see, for example, Kimber-Martin-Ryskin (KMR) [70] and Jung-Salam (JS) [71]), which are not used in our study mostly because of technical difficulties. Note that all above parameterizations give quite similar results excepting, perhaps, the contributions from the small $k_{\perp}^2$ -range: $k_{\perp}^2 \leq 1 \text{ GeV}^2$ (see Ref. [11] and references therein). Because we use $Q_0^2 = 4 \text{ GeV}^2$ in the study of SF F_L , our results depend very slightly on the the small $k_{\perp}^2$ -range of the parameterizations. In the case JB, GBW and KMS sets this observation is supported below by our results and we expect that the application of KMR and JS sets should not strongly change our results.

The differences observed between the curves 2, 3 and 4 are due to the different behavior of the unintegrated gluon distribution as function x and $k_{\perp}$. We see that the SF F_L obtained

in the k_T -factorization approach with KMS and JB parameterizations is close each other ¹⁵ and higher than the SF obtained in the pure perturbative QCD with the GRV gluon density at the leading order approximation. Otherwise, the k_T -factorization approach with GBW parameterization is very close to pure QCD predictions ¹⁶: it should be so because GBW model has deviations from perturbative QCD only at quite low Q^2 values. Thus, the predictions of perturbative QCD and ones based on k_T factorization approach are in agreement each other and with all data within modern experimental uncertainties. So, a possible high values of high-twist corrections to SF F_L predicted in [40] can be important only at low Q^2 values: $Q^2 \leq Q_0^2 = 4 \text{ GeV}^2$.

Fig. 3 is similar to Fig. 2 with one excepting: we add 'experimental data' obtained using the relation between SF $F_L(x, Q^2)$, $F_2(x, Q^2)$ and $dF_2(x, Q^2)/d \ln Q^2$ (see Section 3) as black stars. Because the corresponding data for SF $F_2(x, Q^2)$ and $dF_2(x, Q^2)/d \ln Q^2$ essentially more precise (see [15]) to compare with the preliminary data [18] for F_L , the 'experimental data' have strongly suppressed uncertainties. As it is shown on Fig. 3 there are very good agreement between the new preliminary data [18], the 'experimental data' and predictions of perturbative QCD and k_T -factorization approach.

To estimate of the value of charm mass effect, we recalculate SF F_L^c also in massless approximation similar to (32). In Fig. 4 we show importance of exact m_c -dependence in hard SF of F_L^c to compare with its massless approximation, where we should have $F_L^c(m_c = 0)/F_L(m_c = 0) = 2/5$. As it is possible to see on Fig. 4, the ratio F_L^c/F_L goes to massless limit 2/5 only at asymptotically large Q^2 values.

5 Conclusions

We have applied in the framework of k_T -factorization approach the results of the calculation of the perturbative parts for the structure functions F_L and F_L^c for a gluon target having nonzero momentum square, in the process of photon-gluon fusion to the analysis of present data for the structure function F_L ¹⁷. The analysis has been performed with several parameterizations of unintegrated gluon distributions, for comparison. We have found good agreement between all existing experimental data, the predictions for F_L obtained from the relation between SF $F_L(x, Q^2)$, $F_2(x, Q^2)$ and $dF_2(x, Q^2)/d \ln Q^2$ and the results obtained in the framework of perturbative QCD and ones based on k_T -factorization approach with the three different parameterizations of unintegrated gluon distributions.

We note that it could be also very useful to evaluate the SF F_2 itself ¹⁸ and the derivatives of F_2 respect to the logarithms of $1/x$ and Q^2 with our expressions using the unintegrated gluons. We are considering to present this work and also the predictions for the ratio $R = \sigma_L/\sigma_T$ in a forthcoming article.

¹⁵Note that very similar results have been obtained also for Ryskin-Shabelsky parameterization [72] (see [73]).

¹⁶This fact is evident also from quite large value of $Q_0^2 = 4 \text{ GeV}^2$ chosen here.

¹⁷In Ref [26] we have also obtained quite large contribution of SF F_L^c at low x and high Q^2 ($Q^2 \geq 30 \text{ GeV}^2$).

¹⁸A study of the SF F_2 in the framework of k_T -factorization has been already done in Ref. [74].

The consideration of the SF F_2 in the framework of the leading-twist approximation of perturbative QCD (i.e. for “pure” perturbative QCD) leads to very good agreement (see Ref. [45] and references therein) with the HERA data at low x and $Q^2 \geq 1.5 \text{ GeV}^2$. The agreement improves at lower Q^2 when higher twist terms are taken into account in Ref. [38, 39]. As it has been studied in Refs. [45, 38], the SF F_2 at low Q^2 is sensitive to the small- x behavior of quark distributions. Thus, our future analysis of F_2 in broad Q^2 range in the framework of k_T -factorization approach should require the incorporation of parameterizations of unintegrated quark densities, introduced recently (see Ref. [70, 11] and references therein).

Acknowledgements

We are grateful to Professor Catani for useful discussions and comments.

The study is supported in part by the RFBR grant 02-02-17513. One of the authors (A.V.K.) is supported in part by Heisenberg-Landau program and INTAS grant N366. N.P.Z. also acknowledge the support of Royal Swedish Academy of Sciences.

References

- [1] A.M.Cooper-Sarkar, G.Ingelman, K.R.Long, R.G.Roberts and D.H.Saxon, *Z.Phys.* **C39** (1988) 281;
- [2] L. Bauerdick, A. Glazov and M.Klein, *in Proc. of the Int. Workshop on Future Physics on HERA*, Hamburg, DESY (1996), p.77 (hep-ex/9609017).
- [3] S. Keller, M. Miramontes, G. Parente, J. Sanchez-Guillen, and O.A. Sampayo, *Phys.Lett.* **B270** (1990) 61;
L.H. Orr and W.J. Stirling, *Phys.Rev.Lett.* **B66** (1991) 1673;
E. Berger and R. Meng, *Phys.Lett.* **B304** (1993) 318.
- [4] A.V. Kotikov, *JETP Lett.* **59** (1994) 1; *Phys. Lett.* **B338** (1994) 349.
- [5] R.S. Thorne, *in: Proc. of the Int. Workshop on Deep Inelastic Scattering* (2002), Cracow.
- [6] Yu.L. Dokshitzer, D.V. Shirkov, *Z. Phys.* **C67** (1995) 449;
- [7] W.K. Wong, *Phys. Rev.* **D54** (1996) 1094.
- [8] S.J. Brodsky, V.S. Fadin, V.T. Kim, L.N. Lipatov and G.B. Pivovarov, *JETP. Lett.* **70** (1999) 155;
V.T. Kim, L.N. Lipatov and G.B. Pivovarov, *in: Proceedings of the VIIIth Blois Workshop at IHEP*, Protvino, Russia, 1999 (IITAP-99-013, hep-ph/9911228); *in: Proceedings of the Symposium on Multiparticle Dynamics (ISMD99)*, Providence, Rhode Island, 1999 (IITAP-99-014, hep-ph/9911242).
- [9] S.J. Brodsky, V.S. Fadin, V.T. Kim, L.N. Lipatov and G.B. Pivovarov, *in: Proceedings of the PHOTON2001*, Ascona, Switzerland, 2001 (CERN-TH/2001-341, SLAC-PUB-9069, hep-ph/0111390);

- [10] M. Ciafaloni, D. Colferai and G.P. Salam, Phys. Rev. **D60** (1999) 114036; JHEP **07** (2000) 054;
R.S. Thorne, Phys. Lett. **B474** (2000) 372; Phys. Rev. **D60** (1999) 054031; **D64** (2001) 074005;
G. Altarelli, R.D. Ball and S. Forte, Nucl. Phys. **B621** (2002) 359.
- [11] Bo Andersson et al., hep-ph/0204115.
- [12] L.N. Lipatov, Sov. J. Nucl. Phys. **23** (1976) 338;
E.A. Kuraev, L.N. Lipatov and V.S. Fadin, Sov. Phys. JETP **44** (1976) 443, **45** (1977) 199;
Ya.Ya. Balitzki and L.N. Lipatov, Sov. J. Nucl. Phys. **28** (1978) 822;
L.N. Lipatov, Sov. Phys. JETP **63** (1986) 904.
- [13] H1 Collab.: S. Aid et al., Phys.Lett. **B393** (1997) 452.
- [14] R.S. Thorne, Phys.Lett. **B418** (1998) 371.
- [15] H1 Collab.: S. Adloff et al., Eur. Phys. J. **C21** (2001) 33
- [16] H1 Collab.: D. Eckstein, *in* Proc. Int. Workshop on Deep Inelastic Scattering, (2001), Bologna;
H1 Collab.: M.Klein, *in* Proc. of the 9th Int. Workshop on Deep Inelastic Scattering, DIS 2001 (2001), Bologna.
- [17] H1 Collab.: T. Lastovicka, *in* Proc. of the 10th Int. Workshop on Deep Inelastic Scattering, DIS 2002 (2002), Cracow;
H1 Collab.: J. Gayler, *in* Proc. of the 10th Int. Workshop on Deep Inelastic Scattering, DIS 2002 (2002), Cracow.
- [18] H1 Collab., N. Gogitidze, J. Phys. **G28** (2002) 751 (hep-ph/0201047)
- [19] A.V. Kotikov, JETP **80** (1995) 979.
- [20] A.V. Kotikov and G. Parente, Mod. Phys. Lett. **A12** (1997) 963; *in* Proc. Int. Workshop on Deep Inelastic Scattering and Related Phenomena (1996), Rome, p. 237 (hep-ph/9608409);
- [21] A.V. Kotikov and G. Parente, JETP **85** (1997) 17; hep-ph/9609439.
- [22] S. Catani, M. Ciafaloni and F. Hautmann, Phys. Lett. **B242** (1990) 97; Nucl. Phys. **B366** (1991) 135; Nucl. Phys. B (Proc.Suppl.) **29A** (1992) 182; Preprint CERN - TH.6398/92, *in* Proceeding of the Workshop on Physics at HERA (Hamburg, 1991), v.2, p.690;
- [23] J.C. Collins and R.K. Ellis, Nucl. Phys. **B360** (1991) 3.
- [24] E.M. Levin, M.G. Ryskin, Yu.M. Shabelskii and A.G. Shuvaev, Sov. J. Nucl. Phys. **53** (1991) 657.

- [25] S. Catani and F. Hautmann, Nucl. Phys. **B427** (1994) 475;
S. Catani, Preprint DFF 254-7-96 (hep-ph/9608310).
- [26] A.V. Kotikov, A. V. Lipatov, G. Parente and N.P. Zotov, Preprint US-FT/7-01 (hep-ph/0107135); *in* Proc. of the XVIth International Workshop “High Energy Physics and Quantum Field Theory” (2001), Moscow.
- [27] V.N. Baier, V.S. Fadin and V.A. Khose, Zh.Eksp.Teor.Fiz. **50** (1966) 156 [Sov. J. JETP **23** (1966) 104];
V.N. Baier, V.M. Katkov and V.S. Fadin, *Relativistic electron radiation*, (Moscow, Atomizdat, 1973) (in Russian);
V.G. Zima, Yad. Fiz. **16** (1972) 1051 [Sov. J. Nucl. Phys. **16** (1973) 580].
- [28] V.M. Budnev, I.F. Ginsburg, G.V. Meledin and V.G. Serbo, Phys. Rept. **15** (1975) 181.
- [29] G. Bottazzi, G. Marchesini, G.P. Salam, and M. Scorletti, JHEP **9812** (1998) 011.
- [30] V.N. Gribov and L.N. Lipatov, Sov. J. Nucl. Phys. **15** (1972) 438, **15** (1972) 675;
L.N. Lipatov, Sov. J. Nucl. Phys. **20** (1975) 94;
G. Altarelli and G. Parisi, Nucl. Phys. **B126** (1977) 298;
Yu.L. Dokshitzer, Sov. Phys. JETP **46** (1977) 641.
- [31] E.A. Kuraev and L.N. Lipatov, Yad. Fiz. **16** (1972) 1060 [Sov. J. Nucl. Phys. **16** (1973) 584].
- [32] K. Prytz, Phys.Lett. **B311** (1993) 286.
- [33] A.V. Kotikov, JETP Lett. **59** (1994) 667;
A.V. Kotikov and G. Parente, Phys.Lett. **B379** (1996) 195.
- [34] H1 Collab., T. Ahmed et al., Nucl. Phys. **B439** (1995) 471.
- [35] ZEUS Collab., M. Derrick et al., Z. Phys. **C65** (1995) 379.
- [36] A.V. Kotikov, Phys. Atom. Nucl. **57** (1994) 133; Phys. Rev. **D49** (1994) 5746.
- [37] F. Martin, Phys. Rev. **D19** (1979) 1382;
C. Lopez and F.I. Yndurain, Nucl. Phys. **B171** (1980) 231; **B183** (1981) 157.
- [38] A.V. Kotikov and G. Parente, *in* Proc. Int. Seminar Relativistic Nuclear Physics and Quantum Chromodynamics (2000), Dubna (hep-ph/0012299); *in* Proc. of the 9th Int. Workshop on Deep Inelastic Scattering, DIS 2001 (2001), Bologna (hep-ph/0106175).
- [39] V.G. Krivokhijine and A.V. Kotikov, JINR preprint E2-2001-190 (hep-ph/0108224); *in*: Proc. of the XVIth International Workshop “High Energy Physics and Quantum Field Theory” (2001), Moscow (hep-ph/0206221); *in*: Proc. of the Int. Workshop “Renormalization Group 2002” (2002), High Tatras, Slovakia (hep-ph/0207222).
- [40] J. Bartels, K.Golec-Biernat, and K. Peters, Eur.Phys.J. **C17** (2000) 121;
J. Bartels, *in*: Proc. of the Int. Workshop on Deep Inelastic Scattering (2002), Cracow.

- [41] H1 Collab.: C. Adloff et al., Phys. Lett. **B520** (2001) 183.
- [42] A.D. Martin, W.S. Stirling and R.G. Roberts, *Phys.Lett.* **B387** (1996) 419.
- [43] M. Gluck, E. Reya and A. Vogt, Eur. Phys. J **C5** (1998) 461.
- [44] A.D. Martin, W.S. Stirling, R.G. Roberts and R.S. Thorne, Eur. Phys. J **C23** (2002) 73;
CTEQ Collab.: J. Pumplin et al., Preprint MSU-HEP-011101 (hep-ph/0201195).
- [45] A.V. Kotikov and G. Parente, Nucl. Phys. **B549** (1999) 242; Nucl. Phys. (Proc. Suppl.) **99A** (2001) 196 (hep-ph/0010352); *in* Proc. of the Int. Conference PQFT98 (1998), Dubna (hep-ph/9810223); *in* Proc. of the 8th Int. Workshop on Deep Inelastic Scattering, DIS 2000 (2000), Liverpool, p. 198 (hep-ph/0006197).
- [46] A.V. Kotikov, Mod. Phys. Lett. A **11** (1996) 103; Phys. At. Nucl. **59** (1996) 2137.
- [47] ZEUS Collab., B. Surrow, talk given at the International Europhysics Conference on High Energy Physics, July 2001 (hep-ph/0201025).
- [48] A.V. Kotikov and G. Parente, Preprint US-FT/3-02 (hep-ph/0207276).
- [49] V.N. Fadin and L.N. Lipatov, Phys. Lett. **B429** (1998) 127;
M. Ciafaloni and G. Camici, Phys. Lett. **B430** (1998) 349.
- [50] A.V. Kotikov and L.N. Lipatov, Nucl. Phys. **B582** (2000) 19; *in: Proc. of the XXXV Winter School*, Repino, S'Peterburg, 2001 (hep-ph/0112346).
- [51] A.V. Kotikov, *Phys. Atom. Nucl.* **56** (1993) 1276.
- [52] V.I. Vovk, A.V. Kotikov, and S.I. Maximov, Theor. Math. Phys. **84** (1990) 744;
A.V. Kotikov, S.I. Maximov, and I.S. Parobij, Theor. Math. Phys. **111** (1997) 442.
- [53] L. L. Jenkovszky, A. V. Kotikov and F. Paccanoni, Sov. J. Nucl. Phys. **55** (1992) 1224;
JETP Lett. **58** (1993) 163; Phys. Lett. B **314**(1993) 421.
- [54] V.A. Matveev, R.M. Muradian and A.N. Tavkhelidze, Lett. Nuovo Cim. **7** (1973) 719;
S.J. Brodsky and G.R. Farrar, Phys. Rev. Lett. **31** (1973) 1153;
S.J. Brodsky, J. Ellis, E. Cardi, M. Karliner and M.A. Samuel, Phys. Rev. **D56** (1997) 6980.
- [55] E.G. Floratos, C. Kounnas, and R. Lacage, Nucl. Phys. **B192** (1981) 417.
- [56] D.I. Kazakov and A.V. Kotikov, Theor.Math.Phys. **73** (1987) 1264; Nucl.Phys. **B307** (1988) 721; E: **B345** (1990) 299.
- [57] D.I. Kazakov and A.V. Kotikov, Phys.Lett. **B291** (1992) 171.
- [58] NM Collab., M. Arneodo et al., Nucl. Phys. **B483** (1997) 3; A.J. Milsztajn, *in* Proc. of the 4th Int. Workshop on Deep Inelastic Scattering, DIS96 (1996), Rome, p.220.

- [59] U.K. Yang et al., J. Phys. **G22** (1996) 775;
A. Bodek, *in* Proc. of the 4th Int. Workshop on Deep Inelastic Scattering, DIS96 (1996), Rome, p.213;
A. Bodek, S. Rock, and U.K. Yang, Univ. Rochester preprint, UR-1355, 1995.
- [60] CCFR/NuTeV Collab.: U.K. Yang et al., Phys.Rev.Lett. **87** (2001) 251802;
CCFR/NuTeV Collab.: A. Bodek, *in* Proc. of the 9th Int. Workshop on Deep Inelastic Scattering, DIS 2001 (2001), Bologna (hep-ex/00105067).
- [61] BCDMS Collab., A.C. Benvenuti et al., *Phys. Lett.* **B223** (1989) 485; **B237** (1990) 592.
- [62] SLAC Collab., L.W. Whitlow et al., *Phys. Lett.* **B250** (1990) 193.
- [63] E140X Collab., L.H. Tao et al., *Z. Phys.* **C70** (1996) 387.
- [64] EM Collab., J.J. Aubert et al., *Nucl. Phys.* **B259** (1985) 189; **B293** (1987) 740.
- [65] CDHSW Collab., P. Berge et al., *Z. Phys.* **C49** (1991) 187.
- [66] M.G. Ryskin and Yu.M. Shabelski, *Z. Phys.* **C61** (1994) 517; **C66** (1995) 151.
- [67] J. Kwiecinski, A.D. Martin and A. Stasto, *Phys.Rev.* **D56** (1997) 3991.
- [68] J. Blumlein, Preprints DESY 95-121 (hep-ph/9506403), DESY 95-125 (hep-ph/9506446).
- [69] K. Golec-Biernat and M Wusthoff, *Phys.Rev.* **D59** (1999) 014017; **D60** (1999) 014015, 114023.
- [70] M.A. Kimber, A.D. Martin, and M.G. Ryskin, *Phys. Rev.* **D63** (2001) 114027.
- [71] H. Jung and G. Salam, *Eur.Phys.J.* **C19** (2001) 351; H. Jung, hep-ph/9908497
- [72] M.G. Ryskin and Yu.M. Shabelski, *Z. Phys.* **C61** (1994) 517; **C66** (1995) 151.
- [73] A.V. Kotikov, A. V. Lipatov, G. Parente and N.P. Zotov, *in* Proc. of the International School “Heavy Quark Physics“ (2002), Dubna.
- [74] H. Jung, *Nucl. Phys. (Proc. Suppl.)* **79** (1999) 429.

Figure captions

Fig. 1 The diagrams contributing to $T_{\mu\nu}$ for a gluon target. They should be multiplied by a factor of 2 because of the opposite direction of the fermion loop. The diagram (a) should be also doubled because of crossing symmetry.

Fig. 2 The structure function $F_L(x, Q^2)$ as a function of x for different values of Q^2 compared to experimental data. The H1 data: the first 1997 ones [13], new 2001 ones [15] and preliminary ones [18] are shown as black triangles, circles and squares, respectively. The data of NM [58], CCFR [60] and BCDMS [61] Collaborations are shown as white triangles, circles and squares, respectively. Curves 1, 2, 3 and 4 correspond to SF obtained in the perturbative QCD with the GRV [43] gluon density at the leading order approximation and to SF obtained in the k_T factorization approach with JB (at $Q_0^2 = 4 \text{ GeV}^2$) [68], Kwiecinski-Martin-Stasto (KMS) and GBW [69] parametrizations of unintegrated gluon distribution.

Fig. 3 The structure function $F_L(x, Q^2)$ as a function of x for different values of Q^2 . To compare with Fig. 2 'experimental data' (see [19]-[21] and Section 3) are added as black stars. The 'experimental data' values depend mostly on the derivative $dF_2(x, Q^2)/d \ln Q^2$, which data are known at little different Q^2 values (see [15]). So, the 'experimental data' obtained at 12, 15, 20, 25 and 35 GeV^2 are presented here at 13.4, 15.3, 22.4, 29.6 and 39.7 GeV^2 , respectively.

Fig. 4 The ratio $F_L^c(x, Q^2)/F_L(x, Q^2)$ as a function of x for different values of Q^2 .





Figure 1:

